

## XIAP

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:100 IF 1:50 - 1:100

**Calculated MW:**57kDa

**Observed MW:**Refer to figures

**Immunogen:**

Recombinant protein of human XIAP

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

bhl

**Synonym:**

API3; ILP1; MIHA; XLP2; BIRC4; IAP-3; hIAP3; hIAP-3;

**Catalog #:**A6869

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**331

**Isotype:**IgG

**Swiss Prot:**P98170

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes a protein that belongs to a family of apoptotic suppressor proteins. Members of this family share a conserved motif termed, baculovirus IAP repeat, which is necessary for their anti-apoptotic function. This protein functions through binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2 and inhibits apoptosis induced by menadione, a potent inducer of free radicals, and interleukin 1-beta converting enzyme. This protein also inhibits at least two members of the caspase family of cell-death proteases, caspase-3 and caspase-7. Mutations in this gene are the cause of X-linked lymphoproliferative syndrome. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 2 and 11.

*To place an order, please [Click HERE](#).*