

## DR1

---

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:**19kDa

**Observed MW:**Refer to figures

**Immunogen:**

Recombinant protein of human DR1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

NC2; NC2-BETA;

**Background:**

This gene encodes a TBP- (TATA box-binding protein) associated phosphoprotein that represses both basal and activated levels of transcription. The encoded protein is phosphorylated in vivo and this phosphorylation affects its interaction with TBP. This protein contains a histone fold motif at the amino terminus, a TBP-binding domain, and a glutamine- and alanine-rich region. The binding of DR1 repressor complexes to TBP-promoter complexes may establish a mechanism in which an altered DNA conformation, together with the formation of higher order complexes, inhibits the assembly of the preinitiation complex and controls the rate of RNA polymerase II transcription.

*To place an order, please [Click HERE](#).*

**Catalog #:**A6903

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**1810

**Isotype:**IgG

**Swiss Prot:**Q01658

**Purity:**Affinity purification

For research use only.