UPF2

Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:148kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human UPF2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

HUPF2; RENT2; smg-3;

Polyclonal Antibody

Species: Rabbit

Gene ID:26019

Isotype:IgG

Swiss Prot:Q9HAU5

Purity: Affinity purification

For research use only.

Background:

This gene encodes a protein that is part of a post-splicing multiprotein complex involved in both mRNA nuclear export and mRNA surveillance. mRNA surveillance detects exported mRNAs with truncated open reading frames and initiates nonsense-mediated mRNA decay (NMD). When translation ends upstream from the last exon-exon junction, this triggers NMD to degrade mRNAs containing premature stop codons. This protein is located in the perinuclear area. It interacts with translation release factors and the proteins that are functional homologs of yeast Upf1p and Upf3p. Two splice variants have been found for this gene; both variants encode the same protein.

To place an order, please Click HERE.





