

## ELAC2

**Reactivity:** Human

**Tested applications:** WB IHC IF

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:100

**Calculated MW:** 92kDa

**Observed MW:** Refer to figures

**Immunogen:**

Recombinant protein of human ELAC2

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

ELC2; HPC2; COXPD17;

**Catalog #:** A7128

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 60528

**Isotype:** IgG

**Swiss Prot:** Q9BQ52

**Purity:** Affinity purification

For research use only.

**Background:**

The protein encoded by this gene has a C-terminal domain with tRNA 3' trailer processing endoribonuclease activity, which catalyzes the removal of the 3' trailer from precursor tRNAs. The protein also interacts with activated Smad family member 2 (Smad2) and its nuclear partner forkhead box H1 (also known as FAST-1), and reduced expression can suppress transforming growth factor-beta induced growth arrest. Mutations in this gene result in an increased risk of prostate cancer. Multiple transcript variants encoding different isoforms have been found for this gene.

**To place an order, please [Click HERE](#).**