

## Acetyl-Histone H3-K27

**Reactivity:** Human Mouse Rat Other (Wide Range)

**Tested applications:** WB IHC IF IP CHIP CHIPseq

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200 IP 1:50 - 1:200

ChIP 1:20 - 1:100 CHIPseq 1:20 - 1:100

**Calculated MW:** 16kDa

**Observed MW:** Refer to figures

**Immunogen:**

A synthetic acetylated peptide corresponding to residues surrounding K27 of human Histone H3

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

q

**Synonym:**

H3K27ac; H3t; H3.4; H3/g; H3FT;

**Catalog #:** A7253

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 8290

**Isotype:** IgG

**Swiss Prot:** Q16695

**Purity:** Affinity purification

For research use only.

**Background:**

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

*To place an order, please [Click HERE](#).*