www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

MPI

Reactivity: Human Rat

Tested applications: WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:100

Calculated MW:47kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human MPI

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

PMI; PMI1; CDG1B;

Background:

Phosphomannose isomerase catalyzes the interconversion of fructose-6-phosphate and mannose-6-phosphate and plays a critical role in maintaining the supply of D-mannose derivatives, which are required for most glycosylation reactions. Mutations in the MPI gene were found in patients with carbohydrate-deficient glycoprotein syndrome, type lb. Alternative splicing results in multiple transcript variants.

To place an order, please Click HERE.

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:4351

Isotype:IgG

Swiss Prot:P34949

Purity: Affinity purification

For research use only.





