

STIM1

Reactivity:Human

Tested applications:WB

Recommended Dilution:WB 1:200 - 1:2000

Calculated MW:77kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human STIM1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

GOK; TAM; TAM1; IMD10; STRMK; D11S4896E;

Background:

This gene encodes a type 1 transmembrane protein that mediates Ca²⁺ influx after depletion of intracellular Ca²⁺ stores by gating of store-operated Ca²⁺ influx channels (SOCs). It is one of several genes located in the imprinted gene domain of 11p15.5, an important tumor-suppressor gene region. Alterations in this region have been associated with the Beckwith-Wiedemann syndrome, Wilms tumor, rhabdomyosarcoma, adrenocortical carcinoma, and lung, ovarian, and breast cancer. This gene may play a role in malignancies and disease that involve this region, as well as early hematopoiesis, by mediating attachment to stromal cells. Mutations in this gene are associated with fatal classic Kaposi sarcoma, immunodeficiency due to defects in store-operated calcium entry (SOCE) in fibroblasts, ectodermal dysplasia and tubular aggregate myopathy. This gene is oriented in a head-to-tail configuration with the ribonucleotide reductase 1 gene (RRM1), with the 3' end of this gene situated 1.6 kb from the 5' end of the RRM1 gene. Alternative splicing of this gene results in multiple transcript variants.

To place an order, please [Click HERE](#).

Catalog #:A7411

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:6786

Isotype:IgG

Swiss Prot:Q13586

Purity:Affinity purification

For research use only.