NDUFV2

Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:27kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human NDUFV2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

CI-24k;

Polyclonal Antibody

Species:Rabbit

Gene ID:4729

Isotype:IgG

Swiss Prot:P19404

Purity: Affinity purification

For research use only.

Background:

The NADH-ubiquinone oxidoreductase complex (complex I) of the mitochondrial respiratory chain catalyzes the transfer of electrons from NADH to ubiquinone, and consists of at least 43 subunits. The complex is located in the inner mitochondrial membrane. This gene encodes the 24 kDa subunit of complex I, and is involved in electron transfer. Mutations in this gene are implicated in Parkinson's disease, bipolar disorder, schizophrenia, and have been found in one case of early onset hypertrophic cardiomyopathy and encephalopathy. A non-transcribed pseudogene of this locus is found on chromosome 19.

To place an order, please Click HERE.





