

MAP2K3

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:39kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human MAP2K3

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

iors

Synonym:

MEK3; MKK3; MAPKK3; PRKMK3; SAPKK2; SAPKK-2;

Catalog #:A7718

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:5606

Isotype:IgG

Swiss Prot:P46734

Purity:Affinity purification

For research use only.

Background:

The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is activated by mitogenic and environmental stress, and participates in the MAP kinase-mediated signaling cascade. It phosphorylates and thus activates MAPK14/p38-MAPK. This kinase can be activated by insulin, and is necessary for the expression of glucose transporter. Expression of RAS oncogene is found to result in the accumulation of the active form of this kinase, which thus leads to the constitutive activation of MAPK14, and confers oncogenic transformation of primary cells. The inhibition of this kinase is involved in the pathogenesis of Yersina pseudotuberculosis. Multiple alternatively spliced transcript variants that encode distinct isoforms have been reported for this gene.

To place an order, please [Click HERE](#).