www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

NAT8

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:100 Calculated MW:26kDa Observed MW:Refer to figures Immunogen: Recombinant protein of human NAT8 Storage Buffer: Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

GLA; CML1; CCNAT; Hcml1; ATase2; TSC501; TSC510;

Background:

This gene, isolated using the differential display method to detect tissue-specific genes, is specifically expressed in kidney and liver. The encoded protein shows amino acid sequence similarity to N-acetyltransferases. A similar protein in Xenopus affects cell adhesion and gastrulation movements, and may be localized in the secretory pathway. A highly similar paralog is found in a cluster with this gene.

To place an order, please Click HERE.



Catalog #:A7759 Antibody Type: Polyclonal Antibody Species:Rabbit Gene ID:9027 Isotype:IgG Swiss Prot:Q9UHE5 Purity:Affinity purification

For research use only.



