For research use only.



CHRFAM7A

Reactivity: Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:46kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human CHRFAM7A

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

D-10; CHRNA7; CHRNA7-DR1;

Background:

The nicotinic acetylcholine receptors (nAChRs) are members of a superfamily of ligand-gated ion channels that mediate fast signal transmission at synapses. The family member CHRNA7, which is located on chromosome 15 in a region associated with several neuropsychiatric disorders, is partially duplicated and forms a hybrid with a novel gene from the family with sequence similarity 7 (FAM7A). Alternative splicing has been observed, and two variants exist, for this hybrid gene. The N-terminally truncated products predicted by the largest open reading frames for each variant would lack the majority of the neurotransmitter-gated ion-channel ligand binding domain but retain the transmembrane region that forms the ion channel. Although current evidence supports transcription of this hybrid gene, translation of the nicotinic acetylcholine receptor-like protein-encoding open reading frames has not been confirmed.

To place an order, please Click HERE.



