

## MRPS22

**Reactivity:** Human Mouse

**Tested applications:** WB IHC

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:100

**Calculated MW:** 41kDa

**Observed MW:** Refer to figures

**Immunogen:**

Recombinant protein of human MRPS22

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

GIBT; GK002; C3orf5; COXPD5; RPMS22; MRP-S22;

**Catalog #:** A8319

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 56945

**Isotype:** IgG

**Swiss Prot:** P82650

**Purity:** Affinity purification

For research use only.

**Background:**

Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein that does not seem to have a counterpart in prokaryotic and fungal-mitochondrial ribosomes. This gene lies telomeric of and is transcribed in the opposite direction from the forkhead box L2 gene. A pseudogene corresponding to this gene is found on chromosome Xq.

**To place an order, please [Click HERE](#).**