www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom

ANAPC10

Reactivity: Human Mouse

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:100

Calculated MW:21kDa

Observed MW: Refer to figures

Immunogen:

Recombinant protein of human ANAPC10

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

DOC1; APC10;

Catalog #:A8330

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID:10393

Isotype: IgG

Swiss Prot:Q9UM13

Purity: Affinity purification

For research use only.

Background:

ANAPC10 is a core subunit of the anaphase-promoting complex (APC), or cyclosome, a ubiquitin protein ligase that is essential for progression through the cell cycle. APC initiates sister chromatid separation by ubiquitinating the anaphase inhibitor securin (PTTG1; MIM 604147) and triggers exit from mitosis by ubiquitinating cyclin B (CCNB1; MIM 123836), the activating subunit of cyclin-dependent kinase-1 (CDK1; MIM 116940) (summary by Wendt et al., 2001

To place an order, please Click HERE.

© 2015 NeoScientific