

MDC1

Reactivity:Human

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:100 - 1:200

Calculated MW:227kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human MDC1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

NFBD1;

Catalog #:A8358

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:9656

Isotype:IgG

Swiss Prot:Q14676

Purity:Affinity purification

For research use only.

Background:

The protein encoded by this gene contains an N-terminal forkhead domain, two BRCA1 C-terminal (BRCT) motifs and a central domain with 13 repetitions of an approximately 41-amino acid sequence. The encoded protein is required to activate the intra-S phase and G2/M phase cell cycle checkpoints in response to DNA damage. This nuclear protein interacts with phosphorylated histone H2AX near sites of DNA double-strand breaks through its BRCT motifs, and facilitates recruitment of the ATM kinase and meiotic recombination 11 protein complex to DNA damage foci.

To place an order, please [Click HERE](#).