TADA2A

Reactivity: Human Mouse Rat

Tested applications:WB

Recommended Dilution: WB 1:500 - 1:2000

Calculated MW:52kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human TADA2A

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

ADA2; ADA2A; KL04P; hADA2; TADA2L;

Polyclonal Antibody

Species: Rabbit

Gene ID:6871

Isotype:IgG

Swiss Prot: 075478

Purity: Affinity purification

For research use only.

Background:

Many DNA-binding transcriptional activator proteins enhance the initiation rate of RNA polymerase II-mediated gene transcription by interacting functionally with the general transcription machinery bound at the basal promoter. Adaptor proteins are usually required for this activation, possibly to acetylate and destabilize nucleosomes, thereby relieving chromatin constraints at the promoter. The protein encoded by this gene is a transcriptional activator adaptor and has been found to be part of the PCAF histone acetylase complex. Several alternatively spliced transcript variants encoding different isoforms of this gene have been described, but the full-length nature of some of these variants has not been determined.

To place an order, please Click HERE.





