

## RYK

**Reactivity:** Human Mouse Rat

**Tested applications:** WB IHC IF

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:20 - 1:200 IF 1:20 - 1:100

**Calculated MW:** 68kDa

**Observed MW:** Refer to figures

**Immunogen:**

A synthetic Peptide of human RYK

**Storage Buffer:**

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Catalog #:** A9502

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 6259

**Isotype:** IgG

**Swiss Prot:** P34925

**Purity:** Affinity purification

For research use only.

**Background:**

The protein encoded by this gene is an atypical member of the family of growth factor receptor protein tyrosine kinases, differing from other members at a number of conserved residues in the activation and nucleotide binding domains. This gene product belongs to a subfamily whose members do not appear to be regulated by phosphorylation in the activation segment. It has been suggested that mediation of biological activity by recruitment of a signaling-competent auxiliary protein may occur through an as yet uncharacterized mechanism. The encoded protein has a leucine-rich extracellular domain with a WIF-type Wnt binding region, a single transmembrane domain, and an intracellular tyrosine kinase domain. This protein is involved in stimulating Wnt signaling pathways such as the regulation of axon pathfinding. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Feb 2012]

**To place an order, please [Click HERE](#).**