## CSE1L

Reactivity: Human

Tested applications:WB

Recommended Dilution: WB 1:500 - 1:2000

Calculated MW:110kDa

Observed MW:Refer to figures

Immunogen:

A synthetic peptide of human CSE1L

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

For research use only.

Species: Rabbit

Gene ID:1434

Swiss Prot:P55060

Purity: Affinity purification

Isotype:IgG

Background:

Proteins that carry a nuclear localization signal (NLS) are transported into the nucleus by the importin-alpha/beta heterodimer. Importin-alpha binds the NLS, while importin-beta mediates translocation through the nuclear pore complex. After translocation, RanGTP binds importin-beta and displaces importin-alpha. Importin-alpha must then be returned to the cytoplasm, leaving the NLS protein behind. The protein encoded by this gene binds strongly to NLS-free importin-alpha, and this binding is released in the cytoplasm by the combined action of RANBP1 and RANGAP1. In addition, the encoded protein may play a role both in apoptosis and in cell proliferation. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jan 2012]

To place an order, please Click HERE.



