

## WDR33

**Reactivity:** Human

**Tested applications:** WB IP

**Recommended Dilution:** WB 1:500 - 1:2000 IP 1:20 - 1:50

**Calculated MW:** 146kDa

**Observed MW:** Refer to figures

**Immunogen:**

Recombinant protein of human WDR33

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Catalog #:** A9531

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 55339

**Isotype:** IgG

**Swiss Prot:** Q9C0J8

**Purity:** Affinity purification

For research use only.

**Background:**

This gene encodes a member of the WD repeat protein family. WD repeats are minimally conserved regions of approximately 40 amino acids typically bracketed by gly-his and trp-aspartate (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene regulation. This gene is highly expressed in testis and the protein is localized to the nucleus. This gene may play important roles in the mechanisms of cytodifferentiation and/or DNA recombination. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

**To place an order, please [Click HERE](#).**