

## Phospho-RB-S795

**Reactivity:** Human

**Tested applications:** WB IF

**Recommended Dilution:** WB 1:500 - 1:1000 IF 1:50 - 1:100

**Calculated MW:** 102kDa

**Observed MW:** Refer to Figures

**Immunogen:**

A phospho specific peptide corresponding to residues surrounding S795 of human RB

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

fk

**Synonym:**

RB1; RB; pRb; OSRC; pp110; p105-Rb

**Catalog #:** AP0088

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 5925

**Isotype:** IgG

**Swiss Prot:** P06400

**Purity:** Affinity purification

For research use only.

**Background:**

The retinoblastoma tumor suppressor protein, Rb, regulates cell proliferation by controlling progression through the restriction point within the G1-phase of the cell cycle (1). Rb has three functionally distinct binding domains and interacts with critical regulatory proteins including the E2F family of transcription factors, c-Abl tyrosine kinase, and proteins with a conserved LXCXE motif (2-4). Cell cycle-dependent phosphorylation by a CDK inhibits Rb target binding and allows cell cycle progression (5). Rb inactivation and subsequent cell cycle progression likely requires an initial phosphorylation by cyclin D-CDK4/6 followed by cyclin E-CDK2 phosphorylation (6). Specificity of different CDK/cyclin complexes has been observed in vitro (6-8) and cyclin D1 is required for Ser780 phosphorylation in vivo (9).

**To place an order, please [Click HERE](#).**