www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom

Phospho-SMC1A-S957

Reactivity: Human

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:20 - 1:100

Calculated MW:143kDa

Observed MW: Refer to Figures

Immunogen:

A phospho specific peptide corresponding to residues surrounding S957 of human SMC1A

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

bf

Synonym:

SMC1; SMCB; CDLS2; SB1.8; SMC1L1; DXS423E; SMC1alpha

Catalog #:AP0090

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID:8243

Isotype: IgG

Swiss Prot:Q14683

Purity: Affinity purification

For research use only.

Background:

Proper cohesion of sister chromatids is a prerequisite for the correct segregation of chromosomes during cell division. The cohesin multiprotein complex is required for sister chromatid cohesion. This complex is composed partly of two structural maintenance of chromosomes (SMC) proteins, SMC3 and either SMC1B or the protein encoded by this gene. Most of the cohesin complexes dissociate from the chromosomes before mitosis, although those complexes at the kinetochore remain. Therefore, the encoded protein is thought to be an important part of functional kinetochores. In addition, this protein interacts with BRCA1 and is phosphorylated by ATM, indicating a potential role for this protein in DNA repair. This gene, which belongs to the SMC gene family, is located in an area of the X-chromosome that escapes X inactivation. Mutations in this gene result in Cornelia de Lange syndrome. Alternative splicing results in multiple transcript variants encoding different isoforms.

To place an order, please Click HERE.

© 2015 NeoScientific

Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)

Request a CATALOG CLICK HERE