

## Phospho-MYC-S373

**Reactivity:** Human

**Tested applications:** WB

**Recommended Dilution:** WB 1:500 - 1:1000

**Calculated MW:** 60kDa

**Observed MW:** Refer to Figures

**Immunogen:**

A phospho specific peptide corresponding to residues surrounding S373 of human MYC

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

bkmoq

**Synonym:**

MRTL; MYCC; c-Myc; bHLHe39;

**Catalog #:** AP0132

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 4609

**Isotype:** IgG

**Swiss Prot:** P01106

**Purity:** Affinity purification

For research use only.

**Background:**

The protein encoded by this gene is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. There is evidence to show that alternative translation initiations from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site result in the production of two isoforms with distinct N-termini. The synthesis of non-AUG initiated protein is suppressed in Burkitt's lymphomas, suggesting its importance in the normal function of this gene.

**To place an order, please [Click HERE](#).**