

Phospho-PLCG1-Y783

Reactivity: Human Mouse Rat

Tested applications: WB

Recommended Dilution: WB 1:500 - 1:2000

Calculated MW: 155kDa

Observed MW: Refer to Figures

Immunogen:

A phospho specific peptide corresponding to residues surrounding Y783 of human PLCG1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

q

Synonym:

PLC1; NCKAP3; PLC-II; PLC148; PLCgamma1;

Catalog #: AP0164

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 5335

Isotype: IgG

Swiss Prot: P19174

Purity: Affinity purification

For research use only.

Background:

The protein encoded by this gene catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for this gene.

To place an order, please [Click HERE](#).