

## Phospho-VASP-S157

**Reactivity:** Human Mouse Rat

**Tested applications:** WB IHC

**Recommended Dilution:** WB 1:500 - 1:1000 IHC 1:50 - 1:100

**Calculated MW:** 50kDa

**Observed MW:** Refer to Figures

**Immunogen:**

A phospho specific peptide corresponding to residues surrounding S157 of human VASP

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

VASP

**Catalog #:** AP0213

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 7408

**Isotype:** IgG

**Swiss Prot:** P50552

**Purity:** Affinity purification

For research use only.

**Background:**

Vasodilator-stimulated phosphoprotein (VASP) is a member of the Ena-VASP protein family. Ena-VASP family members contain an EHV1 N-terminal domain that binds proteins containing E/DFPPPPXD/E motifs and targets Ena-VASP proteins to focal adhesions. In the mid-region of the protein, family members have a proline-rich domain that binds SH3 and WW domain-containing proteins. Their C-terminal EVH2 domain mediates tetramerization and binds both G and F actin. VASP is associated with filamentous actin formation and likely plays a widespread role in cell adhesion and motility. VASP may also be involved in the intracellular signaling pathways that regulate integrin-extracellular matrix interactions. VASP is regulated by the cyclic nucleotide-dependent kinases PKA and PKG.

**To place an order, please [Click HERE](#).**