

Phospho-PTK2B-Y402

Reactivity: Human Mouse Rat

Tested applications: WB IHC IF

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:100 IF 1:100 - 1:200

Calculated MW: 116kDa

Observed MW: Refer to Figures

Immunogen:

A phospho specific peptide corresponding to residues surrounding Y402 of human PTK2B

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

bq

Synonym:

PKB; PTK; CAKB; FAK2; PYK2; CADTK; FADK2; RAFTK;

Catalog #: AP0214

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 2185

Isotype: IgG

Swiss Prot: Q14289

Purity: Affinity purification

For research use only.

Background:

This gene encodes a cytoplasmic protein tyrosine kinase which is involved in calcium-induced regulation of ion channels and activation of the map kinase signaling pathway. The encoded protein may represent an important signaling intermediate between neuropeptide-activated receptors or neurotransmitters that increase calcium flux and the downstream signals that regulate neuronal activity. The encoded protein undergoes rapid tyrosine phosphorylation and activation in response to increases in the intracellular calcium concentration, nicotinic acetylcholine receptor activation, membrane depolarization, or protein kinase C activation. This protein has been shown to bind CRK-associated substrate, nephrocystin, GTPase regulator associated with FAK, and the SH2 domain of GRB2. The encoded protein is a member of the FAK subfamily of protein tyrosine kinases but lacks significant sequence similarity to kinases from other subfamilies. Four transcript variants encoding two different isoforms have been found for this gene.

To place an order, please [Click HERE](#).