888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

Reactivity: Human Mouse Rat

Phospho-SNCA-Y136

www.neobiolab.com info@neobiolab.com

Tested applications: WB IF

Recommended Dilution: WB 1:500 - 1:2000 IF 1:100 - 1:200

Calculated MW:18kDa

Observed MW:Refer to Figures

Immunogen:

A phospho specific peptide corresponding to residues surrounding Y136 of human SNCA

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Concentration:

bi

Synonym:

PD1; NACP; PARK1; PARK4;

Background:

Alpha-synuclein is a member of the synuclein family, which also includes beta- and gamma-synuclein. Synucleins are abundantly expressed in the brain and alpha- and beta-synuclein inhibit phospholipase D2 selectively. SNCA may serve to integrate presynaptic signaling and membrane trafficking. Defects in SNCA have been implicated in the pathogenesis of Parkinson disease. SNCA peptides are a major component of amyloid plaques in the brains of patients with Alzheimer's disease. Four alternatively spliced transcripts encoding two different isoforms have been identified for this gene.

To place an order, please Click HERE.





Species: Rabbit

Gene ID:6622

Isotype:IgG

Swiss Prot:P37840

Purity: Affinity purification

For research use only.