www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

Phospho-MAPK8/MAPK9/MAPK10-T183/Y185

Reactivity: Human Mouse Rat

Tested applications:WB IF

Recommended Dilution: WB 1:500 - 1:2000 IF 1:100 - 1:200

Calculated MW:46/54kDa Observed MW:Refer to Figures

Immunogen:

A phospho specific peptide corresponding to residues surrounding T183/Y185of human MAPK8/MAPK9/MAPK10

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

JNK; JNK1; PRKM8; SAPK1; JNK-46; JNK1A2; SAPK1c/JNK2; SAPK; p54a; JNK2A; JNK2B; PRKM9; JNK-55; SAPK1a; JNK2BETA; p54aSAPK; JNK2ALPHA/JNK3; JNK3A; PRKM10;

SAPK1b; p493F12; p54bSAPK

Background:

Responds to activation by environmental stress and pro-inflammatory cytokines by phosphorylating a number of transcription factors, primarily components of AP-1 such as JUN, JDP2 and ATF2 and thus regulates AP-1 transcriptional activity. In T-cells, JNK1 and JNK2 are required for polarized differentiation of T-helper cells into Th1 cells By similarity. Phosphorylates heat shock factor protein 4 (HSF4). /Responds to activation by environmental stress and pro-inflammatory cytokines by phosphorylating a number of transcription factors, primarily components of AP-1 such as c-Jun and ATF2 and thus regulates AP-1 transcriptional activity. In T-cells, JNK1 and JNK2 are required for polarized differentiation of T-helper cells into Th1 cells. JNK2 isoforms display different binding patterns: a-1 and a-2 preferentially bind to c-Jun, whereas beta-1 and beta-2 bind to ATF2. However, there is no correlation between binding and phosphorylation, which is achieved at about the same efficiency by all isoforms. JUNB is not a substrate for JNK2 a-2, and JUND binds only weakly to it. /Responds to activation by environmental stress and pro-inflammatory cytokines by phosphorylating a number of transcription factors, primarily components of AP-1 such as c-Jun and ATF2 and thus regulates AP-1 transcriptional activity. Required for stress-induced neuronal apoptosis and the pathogenesis of glutamate excitotoxicity

To place an order, please Click HERE.

Catalog #:AP0276

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID:5599/5601/5602

Isotype:IgG

Swiss Prot:P45983/P45984/P53779

Purity: Affinity purification

For research use only.





