www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

Phospho-Cdc25b-S149

Reactivity: Mouse Rat

Tested applications:WB

Recommended Dilution: WB 1:500 - 1:2000

Calculated MW:62kDa

Observed MW:Refer to Figures

Immunogen:

A phospho specific peptide corresponding to residues surrounding S149 of human Cdc25b

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

AI604853;

Background:

CDC25B is a member of the CDC25 family of phosphatases. CDC25B activates the cyclin dependent kinase CDC2 by removing two phosphate groups and it is required for entry into mitosis. CDC25B shuttles between the nucleus and the cytoplasm due to nuclear localization and nuclear export signals. The protein is nuclear in the M and G1 phases of the cell cycle and moves to the cytoplasm during S and G2. CDC25B has oncogenic properties, although its role in tumor formation has not been determined. Multiple transcript variants for this gene exist.

To place an order, please Click HERE.

Catalog #:AP0322

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:994 Isotype:IgG

Swiss Prot:P30305

Purity: Affinity purification

For research use only.





