

Phospho-CHEK2-S516

Reactivity: Human

Tested applications: WB

Recommended Dilution: WB 1:500 - 1:2000

Calculated MW: 62kDa

Observed MW: Refer to Figures

Immunogen:

A phospho specific peptide corresponding to residues surrounding S516 of human CHEK2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

e

Synonym:

CDS1; CHK2; LFS2; RAD53; hCds1; HuCds1; PP1425;

Catalog #: AP0331

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 11200

Isotype: IgG

Swiss Prot: O96017

Purity: Affinity purification

For research use only.

Background:

In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. In addition, this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutations in TP53. Also, mutations in this gene are thought to confer a predisposition to sarcomas, breast cancer, and brain tumors. This nuclear protein is a member of the CDS1 subfamily of serine/threonine protein kinases. Several transcript variants encoding different isoforms have been found for this gene.

To place an order, please [Click HERE](#).