

Phospho-HDAC4-S632

Reactivity:Human Mouse Rat

Tested applications:WB

Recommended Dilution:WB 1:500 - 1:2000

Calculated MW:140kDa

Observed MW:Refer to Figures

Immunogen:

A phospho specific peptide corresponding to residues surrounding S632 of human HDAC4

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

fp

Synonym:

HD4; AHO3; BDMR; HDACA; HA6116; HDAC-4; HDAC-A;

Catalog #:AP0359

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:9759

Isotype:IgG

Swiss Prot:P56524

Purity:Affinity purification

For research use only.

Background:

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. This protein does not bind DNA directly, but through transcription factors MEF2C and MEF2D. It seems to interact in a multiprotein complex with RbAp48 and HDAC3.

To place an order, please [Click HERE](#).