www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

Phospho-PLCG2-Y1217

Reactivity: Human Mouse Rat

Tested applications: IHC

Recommended Dilution: IHC 1:50 - 1:100

Calculated MW:150kDa

Observed MW:Refer to Figures

Immunogen:

A phospho specific peptide corresponding to residues surrounding Y1217 of human PLCG2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Concentration:

Synonym:

FCAS3; APLAID; RP11-510J16.3;

Background:

The protein encoded by this gene is a transmembrane signaling enzyme that catalyzes the conversion of 1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate to 1D-myo-inositol 1,4,5-trisphosphate (IP3) and diacylglycerol (DAG), using calcium as a cofactor. IP3 and DAG are second messenger molecules important for transmitting signals from growth factor receptors and immune system receptors across the cell membrane.

To place an order, please Click HERE.

Catalog #:AP0429

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:5336

Isotype:IgG

Swiss Prot:P16885

Purity: Affinity purification

For research use only.





