

## ENA 78 Human (8-78 a.a.)

**Description:** Epithelial Neutrophil-Activating Protein 78 Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 71 amino acids (8-78 a.a.) and having a molecular mass of 7.8kDa. The CXCL5 is purified by proprietary chromatographic techniques.

**Synonyms:** Small inducible cytokine B5, CXCL5, Epithelial-derived neutrophil-activating protein 78, Neutrophil-activating peptide ENA-78, ENA-78(1-78), chemokine (C-X-C motif) ligand 5, SCYB5.

**Source:** Escherichia Coli.

**Physical Appearance:** Sterile Filtered White lyophilized (freeze-dried) powder.

**Amino Acid Sequence:** LRELRCVCLQ TTQGVHPKMI SNLQVFAIGP QCSKVEVVAS  
LKNGKEICLD PEAPFLKKVI QKILDGGNKE N.

**Purity:** Greater than 97.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

**Formulation:**

Lyophilized from a 0.2

**Stability:**

Lyophilized ENA78 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CXCL5 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Solubility:**

It is recommended to reconstitute the lyophilized ENA-78 in sterile 18M-cm H<sub>2</sub>O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

**Introduction:**

Chemokine (C-X-C motif) ligand 5 (CXCL5) is a small cytokine belonging to the CXC chemokine family that is also known as epithelial-derived neutrophil-activating peptide 78 (ENA-78). It is produced following stimulation of cells with the inflammatory cytokines interleukin-1 or tumor necrosis factor-alpha. Expression of CXCL5 has also been observed in eosinophils, and can be inhibited with the type II interferon IFN-. This chemokine stimulates the chemotaxis of neutrophils possesses angiogenic properties. It elicits these effects by interacting with the cell surface chemokine receptor CXCR2. The gene for CXCL5 is encoded on four exons and is located on human chromosome 4 amongst several other CXC chemokine genes. CXCL5 has been implicated in connective tissue remodelling.

**Biological Activity:**

The biological activity was determined by its ability to chemoattract human peripheral blood

www.neobiolab.com  
info@neobiolab.com  
888.754.5670, +1 617.500.7103 United States  
0800.088.5164, +44 020.8123.1558 United Kingdom  
neutrophils using a concentration range of 10.0-100.0 ng/ml.



---

***To place an order, please [Click HERE](#).***

Catalog #:CHPS-272

For research use only.

© 2015 NeoScientific



neobiolab | [Term](#) | [Quote/Order](#) | [Services](#) | [Products](#) | [Support](#) | [Corporate](#) | [Contact Us](#)



Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)

Request a  
**CATALOG**  
[CLICK HERE](#)