

MCP 2 Mouse

Description: Monocyte Chemotactic Protein-2 Mouse Recombinant produced in E.Coli is a non-glycosylated, Polypeptide chain containing 74 amino acids and having a molecular mass of 8507 Dalton. The MCP-2 is purified by proprietary chromatographic techniques.

Synonyms: Small inducible cytokine A8, CCL8, Monocyte chemotactic protein 2, MCP-2, Monocyte chemoattractant protein 2, HC14, chemokine (C-C motif) ligand 8, MCP2, SCYA8, SCYA10, AB023418, 1810063B20Rik.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: GPKAPVTCC FHVLLKLIPL RVLKSYERIN NIQCPMEAVV
FQTKQGMSLC VDPTQKWVSE YMEILDQKSQ ILQP.

Purity: Greater than 97.0% as determined by (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

Lyophilized from a 0.2

Stability:

Lyophilized MCP-2 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CCL8 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized CCL8 in sterile 18M-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

Chemokine (C-C motif) ligand 8 (CCL8) is a small cytokine belonging to the CC chemokine family that was once called monocyte chemotactic protein-2 (MCP-2). The CCL8 protein is produced as a precursor containing 109 amino acids, which is cleaved to produce mature CCL8 containing 75 amino acids. The gene for CCL8 is encoded by 3 exons and is located within a large cluster of CC chemokines on chromosome 17q11.2 in humans. MCP-2 is chemotactic for and activates a many different immune cells, including mast cells, eosinophils and basophils, (that are implicated in allergic responses), and monocytes, T cells, and NK cells that are involved in the inflammatory response. CCL8 elicits its effects by binding to several different cell surface receptors called chemokine receptors. These receptors include CCR1, CCR2B and CCR5.

Biological Activity:

Determined by its ability to chemoattract human peripheral blood monocytes using a concentration

www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom
range of 10-100ng/ml corresponding to a Specific Activity of 10,000-100,000IU/mg.



To place an order, please [Click HERE](#).

Catalog #:CHPS-362

For research use only.

© 2015 NeoScientific



Request a
CATALOG
[CLICK HERE](#)