www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

HGF Human, HEK

Description: HGF Human Recombinant produced in HEK cells is a glycosylated single chain peptide, containing 697 a.a. (Gln-32 to Ser-728) having a total molecular weight of 70kDa. The HGF is purified by proprietary chromatographic techniques.

Catalog #:CYPS-097

For research use only.

Synonyms: Scatter Factor (SF), Hepatopoietin (HPTA), HGF, HGFB, F-TCF.

Source: HEK.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: 32-QRKRRNTIHE FKKSAKTTLI KIDPALKIKT KKVNTADQCA NRCTRNKGLP FTCKAFVFDK ARKQCLWFPF NSMSSGVKKE FGHEFDLYEN KDYIRNCIIG KGRSYKGTVS ITKSGIKCOP WSSMIPHEHS FLPSSYRGKD LOENYCRNPR GEEGGPWCFT SNPEVRYEVC DIPQCSEVEC MTCNGESYRG LMDHTESGKI CQRWDHQTPH RHKFLPERYP **DKGFDDNYCR**

Purity: Greater than 95% as obsereved by SDS-PAGE.

Formulation:

The HGF was lyophilized from 1mg/ml in 1xPBS.

Stability:

Lyophilized HGF although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution HGF should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolabs products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized HGF in sterile PBS containing 0.1% endotoxin-free recombinant HSA.

Introduction:

Hepatocyte Growth Factor (HGF) is a multifunctional growth factor which regulates both cell growth and cell motility. It exerts a strong mitogenic effect on hepatocytes and primary epithelial cells. HGF synergizes with Interleukin-3 and GM-CSF to stimulate colony formation of hematopoietic progenitor cells in vitro and may, therefore, also modulate hematopoiesis.

Biological Activity:

The specific activity was determined by the dose-dependent stimulation of the proliferation of the monkey epithelial cell line 4MBr-5 and is typically 10-50ng/ml.

To place an order, please Click HERE.





