

## IL 3 Human, HEK

**Description:** Interleukin-3 Human Recombinant produced in HEK cells is a glycosylated monomer, having a molecular weight range of 17-45kDa due to glycosylation. The IL-3 is purified by proprietary chromatographic techniques.

**Catalog #:** CYPs-103

For research use only.

**Synonyms:** MCGF (Mast cell growth factor), Multi-CSF, HCGF, P-cell stimulation factor, IL-3, MGC79398, MGC79399.

**Source:** HEK.

**Physical Appearance:** Sterile Filtered White lyophilized (freeze-dried) powder.

**Purity:** Greater than 90% as observed by SDS-PAGE.

**Formulation:**

The IL3 was lyophilized from 1mg/ml in 1xPBS.

**Stability:**

Lyophilized IL-3 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL3 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

**Usage:**

NeoBiolabs products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Solubility:**

It is recommended to reconstitute the lyophilized IL-3 in sterile water not less than 100

**Introduction:**

IL3 is a potent growth promoting cytokine. This cytokine is capable of supporting the proliferation of a broad range of hematopoietic cell types. It is involved in a variety of cell activities such as cell growth, differentiation and apoptosis. This cytokine has been shown to also possess neurotrophic activity, and it may be associated with neurologic disorders.

**Biological Activity:**

The activity was determined by the dose dependent stimulation of the proliferation of human TF-1 cells (human erythroleukemic indicator cell line) and is typically 0.5-2ng/ml corresponding to a specific activity of 500,000-2,000,000units/mg.

**To place an order, please [Click HERE](#).**