

TGF b 2 Human, HEK

Description: TGF-b 2 Human Recombinant produced in HEK cells is a non-glycosylated homodimer, having a total molecular weight of 25kDa. The TGF-b 2 is purified by proprietary chromatographic techniques.

Catalog #: CYPs-119

Synonyms: Transforming growth factor, beta 2, cetermin, Glioblastoma-derived T-cell suppressor factor, polygerin, G-TSF, TGF-beta2, TGF-beta-2, transforming growth factor beta-2, BSC-1 cell growth inhibitor, TGFB-2.

For research use only.

Source: HEK.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Purity: Greater than 95% as observed by SDS-PAGE.

Formulation:

The TGF-b 2 was lyophilized from 1mg/ml in 1xPBS.

Stability:

Lyophilized TGF-b 2 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution TGF-b 2 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolabs products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized TGF-b 2 in sterile water not less than 100

Introduction:

TGFB2 is a 27.08 kDa protein having two identical 118 amino acid peptide chains linked by a single disulfide bond. TGFB2 is part of a family of five related cytokines that have an extensive variation of normal and neoplastic cells, indicating the importance of these homo-dimer proteins as multi-functional regulators of cellular activity. The three mammalian isoforms of TGF-beta (TGFB1, TGFB2 and TGFB3) signal through the same receptor and stimulate similar biological responses. They are involved in physiological processes as embryogenesis, tissue remodelling and wound healing.

Biological Activity:

The specific activity was determined by the dose-dependent inhibition of IL-4 induced proliferation of mouse HT-2 cells (BALB/c spleen activated by sheep erythrocytes in the presence of IL-2) and is typically 0.1-0.5ng/ml.

To place an order, please [Click HERE](#).