www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

IL 4 Human

SCIENTIFIC

Description:Interleukin-4 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 130 amino acids and having a molecular mass of 15000 Dalton. The IL-4 is purified by proprietary chromatographic techniques.

Synonyms:BCGF, BCDF, B cell stimulating factor, BSF-1, Lymphocyte stimulatory factor 1, IL-4, MGC79402, Binetrakin, Pitrakinra.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: The sequence of the first five N-terminal amino acids was determined and was found to be Met-His-Lys-Cys-Asp.

Purity:Greater than 98.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Formulation:

Lyophilized from a 0.2

Stability:

Lyophilized Interleukin-4 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL4 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drµgs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Interleukin-4 in sterile 18M-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

IL4 is a pleiotropic cytokine produced by activated T cells. IL4 is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. IL4, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

Biological Activity:

The ED50 as determined by the dose-dependant stimulation of TF-1 cells is < 0.2 ng/ml, corresponding to a Specific Activity of 5,000,000 IU/mg.





www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

References:

he Journal of Immunology, 2006, 177: 6695-6704.Copyright © 2006 by The American Association of Immunologists, Inc.Link:http://www.jimmunol.org/content/177/10/6695.full2.Title:CD26, adenosine deaminase, and adenosine receptors mediate costimulatory signals in the immunological synapsePublication:Published online before print June 27, 2005, doi: 10.1073/pnas.0501050102 PNAS July 5, 2005 vol. 102 no. 27 9583-9588 Link:http://www.pnas.org/content/102/27/9583.full3.Title:A Network Modeling Approach to Analysis of the Th2 Memory Responses Underlying Human Atopic DiseasePublication:doi: 10.4049/jimmunol.0804125 The Journal of Immunology May 15, 2009 vol. 182 no. 10 6011-6021 Link:http://www.jimmunol.org/content/182/10/6011.full4.Title:Defective IL-10 production in severe phenotypes of Crohns diseasePublication:Published online before print February 23, 2009, doi: 10.1189/jlb.1108698 May 2009 Journal of Leukocyte Biology vol. 85 no. 5 896-903 Link:http://www.jleukbio.org/content/85/5/896.full5.Title:Adenosine deaminase potentiates the generation of effector, memory, and regulatory CD4+ T cells .Publication:Published online before print October 19, 2010, doi: 10.1189/jlb.1009696 January 2011 Journal of Leukocyte Biology vol. 89 no. 1 127-136 .Link:http://www.jleukbio.org/content/89/1/127.full6.Title:Glutamate Released by Dendritic Cells as a Novel Modulator of T Cell Activation/Publication:Link:http://journal.9med.net/qikan/article.phpid=3677937.Title:Airway Epithelial Cells Regulate the Functional Phenotype of Locally Differentiating Dendritic Cells: Implications for the Pathogenesis of Infectious and Allergic Airway Disease.Publication: The Journal of Immunology January 1, 2009 vol. 182 no. 1 72-83 Link:http://www.jimmunol.org/content/182/1/72.full8. Title: Inuence of heat stress on human monocyte-derived dendritic cell functions with immunotherapeutic potential for antitumor vaccines.Publication: Published online before printFebruary 20, 2007, doi:10.1189/jlb.0506347May 2007Journal of Leukocyte Biologyvol. 81no. 51179-1187. Link: http://www.jleukbio.org/content/81/5/1179.full.pdf+html

To place an order, please Click HERE.



Catalog #:CYPS-218

For research use only.



