

IL 6 Human

Description: Interleukin-6 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 184 amino acids and having a molecular mass of 21000 Dalton. The IL6 is purified by proprietary chromatographic techniques.

Synonyms: IFN- β 2, B cell differentiation factor, BCDF, BSF-2, HPGF, HSF, MGI-2, B-cell stimulatory factor 2, Interferon beta-2, Hybridoma growth factor, CTL differentiation factor, CDF, IL-6, HGF.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: The sequence of the first five N-terminal amino acids was determined and was found to be Met-Pro-Val-Pro-Pro.

Purity: Greater than 97.0% as determined by (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

Lyophilized from a 0.2 μ m filtered concentrated (1mg/ml) solution in PBS, pH 7.4.

Stability:

Lyophilized Interleukin-6 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL6 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. They may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Interleukin-6 in sterile 18M-cm H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions.

Introduction:

IL-6 is a cytokine with a wide variety of biological functions: it plays an essential role in the final differentiation of b-cells into ig-secreting cells, it induces myeloma and plasmacytoma growth, it induces nerve cells differentiation, in hepatocytes it induces acute phase reactants.

Biological Activity:

The ED₅₀ as determined by the dose-dependant stimulation of murine 7TD1 cells is less than 0.1 ng/ml, corresponding to the specific activity of 1.0 x 10,000,000 Units per mg.

References:

1. Title: Identification of -Escin as a Novel Inhibitor of Signal Transducer and Activator of Transcription 3/Janus-Activated Kinase 2 Signaling Pathway that Suppresses Proliferation and Induces Apoptosis in Human Hepatocellular Carcinoma Cells .Publication: Published online before

www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom

print April 8, 2010, doi: 10.1124/jpet.110.165498 JPET July 2010 vol. 334 no. 1 285-293

Link:<http://jpet.aspetjournals.org/content/334/1/285.full2>.Title:Suppression of Signal Transducer

and Activator of Transcription 3 Activation by Butein Inhibits Growth of Human Hepatocellular

Carcinoma In Vivo.Publication:Published OnlineFirst December 3, 2010; doi:

10.1158/1078-0432.CCR-10-1123 Clin Cancer Res March 15, 2011 17;

1425.Link:<http://clincancerres.aacrjournals.org/content/17/6/1425.full3>.Title:Identification of -escin

as a novel inhibitor of STAT3/JAK2 signaling pathway that suppresses proliferation and induces

apoptosis in human hepatocellular carcinoma cells.Publication:Published online before print April

8, 2010, doi: 10.1124/jpet.110.165498 JPET April 8, 2010

jpet.110.165498Link:<http://jpet.aspetjournals.org/content/early/2010/04/09/jpet.110.165498.short4>.

Title:enhanced self -renewal of hematopoietic stem/progenitor cells mediated by the stem cell

gene

sall4.Publication:Link:<http://www.biomedcentral.com/content/pdf/1756-8722-4-38.pdf5>.Title:Thymo

quinone inhibits proliferation, induces apoptosis and chemosensitizes human multiple myeloma

cells through suppression of signal transducer and activator of transcription 3 activation

pathway.Publication:Article first published online: 10 MAY 2010DOI:

10.1111/j.1476-5381.2010.00874.xLink:<http://onlinelibrary.wiley.com/doi/10.1111/j.1476-5381.2010.00874.x/full>

0.00874.x/full

To place an order, please Click [HERE](#).



Catalog #:CYP5-220

For research use only.