

## IL 17 Human

**Description:** Interleukin-17A Human Recombinant produced in E.Coli is a homodimeric, non-glycosylated polypeptide chain containing a total of 264 amino acids (2 chains of 132 aa) and having a molecular mass of 31kDa. The IL-17 is purified by proprietary chromatographic techniques.

**Catalog #:** CYP5-257

For research use only.

**Synonyms:** CTLA-8, IL-17, IL-17A, Cytotoxic T-lymphocyte-associated antigen 8.

**Source:** Escherichia Coli.

**Physical Appearance:** Sterile Filtered White lyophilized (freeze-dried) powder.

**Purity:** Greater than 95.0% as determined by SDS-PAGE.

**Formulation:**

Sterile filtered lyophilized (freeze-dried) powder, with 20mM Citrate, 0.1M NaCl, pH4.5.

**Stability:**

Lyophilized Interleukin 17A although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL17 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Solubility:**

It is recommended to reconstitute the lyophilized Interleukin 17 in sterile 18M-cm H<sub>2</sub>O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

**Introduction:**

IL17 is a proinflammatory cytokine produced by activated T cells. IL-17 regulates the activities of NF-kappaB and mitogen-activated protein kinases. Interleukin-17 can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of IL-17 are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis.

**Biological Activity:**

The ED50 as determined by the dose-dependent induction of IL-6 in Hs68 cell line was found to be approximately 2ng/ml corresponding to a Specific Activity of 500,000IU/mg.

**To place an order, please [Click HERE](#).**