

IL 4 Human, CHO

Description: Interleukin-4 Human Recombinant produced in CHO is a single, non-glycosylated polypeptide chain containing 129 amino acids and having a molecular mass of 14 kDa, the glycosylation of IL-4 migrates as 18 kDa on SDS-PAGE. The IL-4 is purified by proprietary chromatographic techniques.

Synonyms: BCGF, BCDF, B cell stimulating factor, BSF-1, Lymphocyte stimulatory factor 1, IL-4, MGC79402, Binetrakin, Pitrakinra.

Source: Chinese Hamster Ovarian Cells.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Purity: Greater than 97.0% as determined by (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

Lyophilized from a Phosphate- Buffered Saline (pH 7.4).

Stability:

Lyophilized Interleukin-4 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL4 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Interleukin-4 in sterile 18M-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

IL4 is a pleiotropic cytokine produced by activated T cells. IL4 is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. IL4, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

Biological Activity:

Measured in a cell proliferation assay using TF-1 cells. The ED50 for this effect is 0.05-0.2ng/ml corresponding to a Specific Activity of 5,000,000-20,000,000IU/mg.

References:

Title: A Network Modeling Approach to Analysis of the Th2 Memory Responses Underlying Human

www.neobiolab.com

info@neobiolab.com

888.754.5670, +1 617.500.7103 United States

0800.088.5164, +44 020.8123.1558 United Kingdom

Atopic DiseasePublication:doi: 10.4049/jimmunol.0804125 The Journal of Immunology May 15,

2009 vol. 182 no. 10 6011-6021 Link:<http://www.jimmunol.org/content/182/10/6011.full>



To place an order, please [Click HERE](#).

Catalog #:CYP5-278

For research use only.

© 2015 NeoScientific



neobiolab | Term | Quote/Order | Services | Products | Support | Corporate | Contact Us

Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)



Request a
CATALOG
[CLICK HERE](#)