

Flt3 Ligand Mouse

Description: Flt3-Ligand Mouse Recombinant produced in E.Coli is non-glycosylated, polypeptide chain containing 163 amino acids and having a molecular mass of 18587 Dalton. Flt3-Ligand is purified by proprietary chromatographic techniques.

Catalog #: CYPs-347

Synonyms: Fms-related tyrosine kinase 3 ligand, FLK2, STK1, CD135, Stem Cell Tyrosine Kinase 1, FLT3LG, Flt3.

For research use only.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: MTPDCYFSHS PISSNFKVKF RELTDHLLKD YPVTVAVNQ
DEKHCKALWS LFLAQRWIEQ LKTVAGSKMQ TLLEDVNTEI HFVTSCTFQP LPECLRFVQT
NISHLLKDTG TQLLALKPCI GKACQNF SRC LEVQCQPDSS TLLPPRSPIA LEATELPEPR PRQ.

Purity: Greater than 98.0% as determined by SDS-PAGE.

Formulation:

The FLT3L Mouse protein was lyophilized without any additives.

Stability:

Lyophilized Flt3-Ligand although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Flt-3 Ligand should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Flt3-Ligand in sterile 18M-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

FLT3 ligand is a receptor for the fl cytokine has a tyrosine-protein kinase activity & a growth factor that regulates proliferation of early hematopoietic cells. Flt3-Ligand synergizes with other CSFs and interleukins to induce growth and differentiation.

Biological Activity:

The ED₅₀, calculated by the dose-dependent proliferation of mouse AML5 cells is 1.3-2.0ng/ml.

To place an order, please Click [HERE](#).