www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom

IL 19 Human

Catalog #:CYPS-370

For research use only.

Description: Interleukin-19 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 155 amino acids and having a molecular mass of 17913 Dalton. The IL-19 is purified by proprietary chromatographic techniques.

Synonyms: Melanoma differentiation association like protein, MDA1, NG.1, ZMDA1, IL-10C, IL-19.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: The sequence of the first five N-terminal amino acids was determined and was found to be Met-Leu-Arg-Arg-Cys.

Purity: Greater than 99.0% as determined by(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Formulation:
Lyophilized from a concentrated (1mg/ml) solution in water containing 10mM sodium citrate pH-5.

Stability:
Lyophilized Interleukin-19 althoμgh stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL19 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.

Usage:
NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drμgs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:
It is recommended to reconstitute the lyophilized Interleukin -19 in sterile 18M-cm H2O at a concentration of 0.5mg/ml or less.

Introduction:
IL19 is a cytokine that belongs to the IL10 cytokine subfamily. IL-19 is found to be preferentially expressed in monocytes. It can bind the IL20 receptor complex and lead to the activation of the signal transducer and activator of transcription 3 (STAT3). A similar cytokine in mouse is reported to up-regulate the expression of IL6 and TNF-alpha and induce apoptosis, which sμggests a role of this cytokine in inflammatory responses. Alternatively spliced transcript variants encoding the distinct isoforms have been described.

Biological Activity:
The activity is determined by it's ability to activate STAT following receptor-ligand interaction.

To place an order, please Click HERE.

© 2015 NeoScientific