

IL 19 Human

Description: Interleukin-19 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 155 amino acids and having a molecular mass of 17913 Dalton. The IL-19 is purified by proprietary chromatographic techniques.

Catalog #: CYPs-370

For research use only.

Synonyms: Melanoma differentiation association like protein, MDA1, NG.1, ZMDA1, IL-10C, IL-19.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: The sequence of the first five N-terminal amino acids was determined and was found to be Met-Leu-Arg-Arg-Cys.

Purity: Greater than 99.0% as determined by (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

Lyophilized from a concentrated (1mg/ml) solution in water containing 10mM sodium citrate pH-5.

Stability:

Lyophilized Interleukin-19 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL19 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Interleukin -19 in sterile 18M-cm H2O at a concentration of 0.5mg/ml or less.

Introduction:

IL19 is a cytokine that belongs to the IL10 cytokine subfamily. IL-19 is found to be preferentially expressed in monocytes. It can bind the IL20 receptor complex and lead to the activation of the signal transducer and activator of transcription 3 (STAT3). A similar cytokine in mouse is reported to up-regulate the expression of IL6 and TNF-alpha and induce apoptosis, which suggests a role of this cytokine in inflammatory responses. Alternatively spliced transcript variants encoding the distinct isoforms have been described.

Biological Activity:

The activity is determined by its ability to activate STAT following receptor-ligand interaction.

To place an order, please [Click HERE](#).