www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom

IL 2 Mouse

Catalog #:CYPS-377

For research use only.

Description: Interleukin-2 Mouse Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 148 amino acids and having a molecular mass of 17231 Dalton. The IL-2 is purified by proprietary chromatographic techniques.

Synonyms: Interleukin-2, T-cell growth factor (TCGF), Aldesleukin, Lymphokine, IL-2.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence:PTSSSTSSST AEAQQQQQQQ QQQQQHLEQL LMDLQELLSR MENYRNLKLP RMLTFKFYLP KQATELKDLQ CLEDELGPLR HVLDLTQSKS FQLEDAENFI SNIRVTVVKL KGSDNTFECQ FDDESATVVD FLRRWIAFCQ SIISTSPQ.

Purity: Greater than 98.0% as determined by(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Formulation:
The protein (1mg/ml) was lyophilized extensive dialysis against 10mM sodium citrate buffer to a pH=4.

Stability:
Lyophilized Interleukin-2 althoμgh stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL2 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.

Usage:
NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drμgs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:
It is recommended to reconstitute the lyophilized Mouse IL-2 in sterile 18M-cm H2O not less than 100μg/ml, which can then be further diluted to other aqueous solutions.

Introduction:
IL2 is a secreted cytokine that is important for the proliferation of T and B lymphocytes. The receptor of this cytokine is a heterotrimeric protein complex whose gamma chain is also shared by interleukin 4 (IL4) and interleukin 7 (IL7). The expression of this gene in mature thymocytes is monoallelic, which represents an unusual regulatory mode for controlling the precise expression of a single gene. The targeted disruption of a similar gene in mice leads to ulcerative colitis-like disease, which sμggests an essential role of this gene in the immune response to antigenic stimuli.

Biological Activity:
The ED50 as determined by the dose-dependant stimulation of murine CTLL-2 cells is 0.18 ng/ml corresponding to a Specific Activity of 55,000,000IU/mg.

To place an order, please Click HERE.

© 2015 NeoScientific

Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)
Request a CATALOG CLICK HERE