

## IFN g Porcine

**Description:** Interferon-gamma Porcine Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 146 amino acids and having a molecular mass of 17140 Dalton. The IFN-gamma is purified by proprietary chromatographic techniques.

Catalog #: CYPs-409

**Synonyms:** Immune Interferon, type II interferon, T cell interferon, MAF, IFNG, IFG, IFI, IFN-gamma.

For research use only.

**Source:** Escherichia Coli.

**Physical Appearance:** Sterile Filtered White lyophilized (freeze-dried) powder.

**Amino Acid Sequence:** The sequence of the first five N-terminal amino acids was determined and was found to be Ser-Tyr-Cys-Gln-Ala.

**Purity:** Greater than 95.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

**Formulation:**

The protein was lyophilized with no additives.

**Stability:**

Lyophilized Interferon-g although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IFN gamma should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Solubility:**

It is recommended to reconstitute the lyophilized Interferon-gamma in sterile 18M-cm H<sub>2</sub>O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

**Introduction:**

IFN-gamma produced by lymphocytes activated by specific antigens or mitogens. IFN-gamma, in addition to having antiviral activity, has important immunoregulatory functions, it is a potent activator of macrophages, and has antiproliferative effects on transformed cells and it can potentiate the antiviral and antitumor effects of the type I interferons.

**Biological Activity:**

The ED<sub>50</sub> as determined by the amount of interferon that inhibited 50% of the cytopathic effect of vesicular stomatitis virus in MDBK cells was found to be 0.2-0.6ng/ml.

**To place an order, please [Click HERE](#).**