

## sCD40 Mouse

**Description:**sCD40 Mouse Recombinant produced in E.Coli is a non-glycosylated, Polypeptide chain containing 149 amino acids and having a molecular mass of 16409 Dalton. The sCD40 is purified by proprietary chromatographic techniques.

**Catalog #:**CYPS-479

For research use only.

**Synonyms:**CD40-L, Tumor necrosis factor ligand superfamily member 5, TNF-related activation protein, TRAP, T cell antigen Gp39, CD154 antigen, sCD40, IGM, IMD3, HIGM1, T-BAM, TNFSF5, hCD40L.

**Source:**Escherichia Coli.

**Physical Appearance:**Sterile Filtered White lyophilized (freeze-dried) powder.

**Amino Acid Sequence:**The sequence of the first five N-terminal amino acids was determined and was found to be Met-Gln-Arg-Gly-Asp.

**Purity:**Greater than 98.0% as determined by(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

**Formulation:**

Lyophilized from a sterile concentrated solution (1mg/ml) with 10mM Sodium Phosphate pH=7.5.

**Stability:**

Lyophilized sCD40 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CD154 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY.They may not be used as drugs,agricultural or pesticidal products, food additives or household chemicals.

**Solubility:**

It is recommended to reconstitute the lyophilized sCD40 in sterile 18M-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

**Introduction:**

CD40L or CD154 is a membrane glycoprotein and differentiation antigen expressed on the surface of T-cells. The CD40 ligand stimulates B-cell proliferation and secretion of all immunoglobulin isotypes in the presence of cytokines . CD40 ligand has been shown to induce cytokine production and tumoricidal activity in peripheral blood monocytes. It also costimulates proliferation of activated T-cells and this is accompanied by the production of IFN-gamma , TNF-alpha , and IL2 .

**Biological Activity:**

The ED50 as determined by its ability to induce MIP-1 alpha & TNF-alpha from mouse splenocytes was found to be 0.1

**To place an order, please [Click HERE](#).**