

## IL31 Mouse

**Description:** IL31 mouse recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 141 amino acids and having a molecular mass of 15.7 kDa.

**Catalog #:** CYPs-611

**Synonyms:** Interleukin 31, IL31, IL-31.

For research use only.

**Source:** Escherichia Coli.

**Physical Appearance:** Sterile Filtered White lyophilized (freeze-dried) powder.

**Amino Acid Sequence:** MTCSLSFGAP ISKEDLRTTI DLLKQESQDL YNNYSIKQAS  
GMSADESIQL PCFSLDREAL TNISVIAHL EKVVLSENT VDTSWVIRWL TNISCFNPLN  
LNISVPGNTD ESYDCKVFVL TVLKQFSNCM AELQAKDNTT C.

**Purity:** Greater than 97.0% as determined by (a) Analysis by SEC-HPLC. (b) Analysis by SDS-PAGE.

**Formulation:**

The IL31 (1mg/ml) was lyophilized from 10mM sodium Phosphate pH-7.4.

**Stability:**

Lyophilized IL31 Recombinant although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL31 should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

**Usage:**

NeoBiolabs products are furnished for LABORATORY RESEARCH USE ONLY. They may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Solubility:**

It is recommended to reconstitute the lyophilized IL31 in sterile 18M-cm H<sub>2</sub>O not less than 100

**Introduction:**

IL-31 produced by activated Th2-type T cells, cooperates with a heterodimeric receptor consisting of IL-31 Receptor Antagonist and Oncostatin-M Receptor that is continuously expressed on epithelial cells and keratinocytes. IL-31 plays a role in the promotion of allergic skin disorders and in regulating other allergic diseases, such as asthma. IL-31 is involved in the itching sensation and endorses the scratching behavior in NC/Nga mice with atopic dermatitis. IL-31 expression is connected with CLA(+) T cells and contributes to the development of atopic dermatitis-induced skin inflammation and pruritus. IL-31 is a powerful inducer of proinflammatory mediators in human colonic SEMFs. IL-31 takes part as a proinflammatory cytokine derived from Th2 cells. Serum IL-31 level is higher in patients with atopic dermatitis. IL-31 is involved in a broad range of immune- & non-immune cells & possesses potential pleiotropic physiological functions, including regulating hematopoiesis & immune re

**To place an order, please [Click HERE](#).**