www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom

IL31 Mouse

Catalog #:CYPS-611

For research use only.

Description:IL31 mouse recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 141 amino acids and having a molecular mass of 15.7 kDa.

Synonyms: Interleukin 31, IL31, IL-31.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence:MTCSLSFGAP ISKEDLRTTI DLLKQESQDL YNNYSIKQAS GMSADESIQL PCFSLDREAL TNISVIIAHL EKVKVLSENT VDTSWVIRWL TNISCFNPLN LNISVPGNTD ESYDCKVFVL TVLKQFSNCM AELQAKDNTT C.

Purity: Greater than 97.0% as determined by(a) Analysis by SEC-HPLC.(b) Analysis by SDS-PAGE.

Formulation:
The IL31 (1mg/ml) was lyophilized from 10mM sodium Phosphate pH-7.4.

Stability:
Lyophilized IL31 Recombinant althoμgh stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL31 should be stored at 4°C between 2-7 days and for future use below -18°C.Please prevent freeze-thaw cycles.

Usage:
NeoBiolabs products are furnished for LABORATORY RESEARCH USE ONLY. They may not be used as drμgs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:
It is recommended to reconstitute the lyophilized IL31 in sterile 18M-cm H2O not less than 100

Introduction:
IL-31 produced by activated Th2-type T cells, cooperates with a heterodimeric receptor consisting of IL-31 Receptor Anatagonist and Onconstatin-M Receptor that is continuesly expressed on epithelial cells and keratinocytes. IL-31 plays a role in the promotion of allergic skin disorders and in regulating other allergic diseases, such as asthma. IL-31 is involved in the itching sensation and endorses the scratching behavior in NC/Nga mice with atopic dermatitis. IL-31 expression is connectd with CLA(+) T cells and contributes to the development of atopic dermatitis-induced skin inflammation and pruritus. IL-31 is a powerful inducer of proinflammatory mediators in human colonic SEMFs. IL-31 takes part as a proinflammatory cytokine derived from Th2 cells.Serum IL-31 level is higher in patients with atopic dermatitis. IL-31 is involved in a broad range of immune- & non-immune cells & possesses potential pleiotropic physiological functions, including regulating hematopoiesis & immune re

To place an order, please Click HERE.

© 2015 NeoScientific

Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)
Request a CATALOG CLICK HERE