

## IL 17A/F Human

**Description:**IL-17A/F Human Recombinant produced in E.Coli is a heterodimeric, non-glycosylated polypeptide chain containing 1 monomeric subunit of each IL-17A & IL-17F. The active dimer contains 271 amino acids and having a total molecular mass of 30.7 kDa. The IL-17A/F Human is purified by proprietary chromatographic techniques.

**Synonyms:**IL17A/F, IL17 A/F, IL-17A/F, IL-17 A/F, IL17AF, IL-17 AF, Interleukin-17 A/F, Interleukin-17 AF.

**Source:**Escherichia Coli.

**Physical Appearance:**Sterile Filtered White lyophilized (freeze-dried) powder.

**Amino Acid Sequence:**MIVKAGITIP RNP GCPNSED KNFPRTVMVN LNIHNRNTNT  
NPKRSSDYNN RSTSPWNLHR NEDPERYPSV IWEAKCRHLG CINADGNVDY HMNSVPIQQE  
ILVLRREPPH CPNSFRLEKI LVSVGCTCVT PIVHHVA.MRKIPKVGHT FFQKPESCPP  
VPGGSMKLDI GIINENQRVS MSRNIESRST SPWNYTVTWD PNRYPSEVVQ AQCRNLGCIN  
AQGKEDISMN SVPIQ

**Purity:**Greater than 98.0% as determined by SDS-PAGE.

**Formulation:**

Lyophilized from a concentrated (1mg/ml) solution containing no additives.

**Stability:**

Lyophilized Human IL-17A/F although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Human IL-17A/F should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Solubility:**

It is recommended to reconstitute the lyophilized Human IL-17A/F in sterile water not more than 1mg/ml, which can then be further diluted to other aqueous solutions.

**Introduction:**

Human IL-17A/F is a 40kDa glycoprotein which is secreted as a disulfide-linked heterodimer. IL-17A/F consists of two proteins of the IL-17 family, IL-17A and IL17F. Proteins of the 6 homodimeric IL17 family show a cysteine knot motif that contains two disulfide-bonds. Human IL17A is produced as a 155 a.a precursor that includes a 23 amino acids signal sequence and a 132 amino acid chain that includes an N-linked glycosylation site. Human IL17F is produced as a 153 amino acid precursor with a 20 amino acid signal sequence and a 133 amino acid region. Similar to IL17A, IL17F also has an N-linked glycosylation site. Both proteins (IL17A & IL17F) share 50% amino acid sequence identity. Human IL17A & IL17F show approximately 60% homology in their amino acid sequence to mouse IL-17A and IL-17F. Interleukin-17A/F and IL17A, IL17F homodimers are manufactured by activated CD4+ T cells, called Th17. IL-23 causes Th17

www.neobiolab.com  
info@neobiolab.com  
888.754.5670, +1 617.500.7103 United States  
0800.088.5164, +44 020.8123.1558 United Kingdom

lymphocytes to manufacture IL-17A/F. IL17RA and IL17RC form a heterodimer for the binding of IL17A and IL17F. IL-17A/F binds IL-17RA. Interleukin-17A/F induces chemokine production and airway neutrophilia with intermediate potency between IL17A (most potent) and IL17F (least potent).

**To place an order, please [Click HERE](#).**



Catalog #:CYP5-630

For research use only.

© 2015 NeoScientific

