

IL 17 A/F Mouse

Description: Interleukin-17 A/F Mouse Recombinant produced in E.Coli is a heterodimeric, non-glycosylated polypeptide comprised of IL17A monomeric subunit & and IL17F monomeric subunit containing a total of 266 amino acids and having a total molecular mass of 29.8kDa. The IL-17 A/F is purified by proprietary chromatographic techniques.

Synonyms: IL17A/F, IL17 A/F, IL-17A/F, IL-17 A/F, IL17AF, IL-17 AF, Interleukin-17 A/F, Interleukin-17 AF.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence:

RKNPKAGVLPALQKAGNCPPLDNTVRVDIRIFNQNGISVPREFQNRSSSPWDYNITRDPHRFPS
EIAEAQCRHSGCINAQGQEDSTMNSVAIQQEILVLRREPQGCSNSFRLEKMLLKVGCTCVKPIVH
QAAAAIIPQSSACPNTAKDFLQNVKVNKVFNSLGAKVSSRRPSDYLNLRSTSPWTLHRNEDPDR
YPSVIWEAQCRHQRCVNAEGKLDHMHNSVLIQQEILVLKREPESCPFTFRVEKMLVGVGC

Purity: Greater than 97.0% as determined by SDS-PAGE.

Formulation:

Lyophilized from a concentrated (1mg/ml) solution containing no additives.

Stability:

Lyophilized Mouse IL17 A/F although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Mouse IL17 A/F should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Mouse IL17 A/F in sterile 18M-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

Human IL-17A/F is a 40kDa glycoprotein which is secreted as a disulfide-linked heterodimer. IL-17A/F consists of two proteins of the IL-17 family, IL-17A and IL17F. Proteins of the 6 homodimeric IL17 family show a cysteine knot motif that contains two disulfide-bonds. Human IL17A is produced as a 155 a.a precursor that includes a 23 amino acids signal sequence and a 132 amino acid chain that includes an N-linked glycosylation site. Human IL17F is produced as a 153 amino acid precursor with a 20 amino acid signal sequence and a 133 amino acid region. Similar to IL17A, IL17F also has an N-linked glycosylation site. Both proteins (IL17A & IL17F) share 50% amino acid sequence identity. Human IL17A & IL17F show approximately 60% homology in their amino acid sequence to mouse IL-17A and IL-17F. Interleukin-17A/F and IL17A, IL17F homodimers are manufactured by activated CD4+ T cells, called Th17. IL-23 causes Th17

www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom

lymphocytes to manufacture IL-17A/F. IL17RA and IL17RC form a heterodimer for the binding of IL17A and IL17F. IL-17A/F binds IL-17RA. Interleukin-17A/F induces chemokine production and airway neutrophilia with intermediate potency between IL17A (most potent) and IL17F (least potent).

To place an order, please [Click HERE](#).



Catalog #:CYP5-647

For research use only.

© 2015 NeoScientific



neobiolab | [Term](#) | [Quote/Order](#) | [Services](#) | [Products](#) | [Support](#) | [Corporate](#) | [Contact Us](#)



Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)

Request a
CATALOG
[CLICK HERE](#)