www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## IL 9 Rat

Description:Interleukin-9 Rat Recombinant produced in E.coli is a single, non-glycosylated polypeptide chain containing 127 amino acids and having a molecular mass of 14.3kDa.The IL-9 is purified by proprietary chromatographic techniques.

Catalog #:CYPS-755

For research use only.

Synonyms:Interleukin 9, Protein II9, II9.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: MQRCSTSWGI QHTSYLIENL KDDPSSKCSC SANVTSCLCL PIPSDDCTTP CFQEGMSQVT NATQQSKFSP FFFRVKRIVE TLKSNKCQFF SCEKPCNQTT AGNTVSFLKS LLKTFQKTEV QVQRSRA.

Purity: Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

#### Formulation:

IL9 protein was lyophilized from a 0.2

#### Stability:

Lyophilized Interleukin-9 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL9 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

# Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

## Solubility:

It is recommended to reconstitute the lyophilized Interleukin 9 in sterile 18M-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

### Introduction:

Factor that is thought to be a regulator of hematopoiesis. It has been shown to enhance the growth of human mast cells and megakaryoblastic leukemic cells as well as murine helper t-cell clones. IL-9 is a glycoprotein with a molecular weight of 32-39 that is derived from T-cells, and maps to human chromosome 5.

#### **Biological Activity:**

The ED50 as determined by a cell proliferation assay using murine TS1 cells is less than 10 ng/ml, corresponding to a specific activity of > 1.0

To place an order, please Click HERE.





